IRVINE PLASTIC SURGERY CENTER

16300 Sand Canyon Avenue, Suite 1011 Irvine, California 92618 Office: (949) 727-3999 Cell: (949) 278-6493

WHAT IS CO2 LASER SKIN RESURFACING?

CO2 is a Carbon Dioxide laser that has been used for more than 25 years in the Aesthetic Industry for treatment of skin conditions such as fine and coarse wrinkles, scars of various origin, uneven pigmentation, and dilated pores. When a CO2 beam of light comes in contact with the epidermis it heats and vaporizes the skin tissue, instantly removing the superficial layers of the skin, scars and wrinkles while smoothing out the surface of the skin. This skin remodeling occurs with new skin and collagen growth.

FRACTIONAL SKIN RESURFACING and its BENEFITS

Fractional Skin Resurfacing is a Revolutionary Delivery System that increases the possibilities for busy people who want healthy, younger looking skin with minimal pain in a short period of time. The MiXto combines the benefits of a CO2 laser with a micro spot size of 300 microns that can treat 20-100% of the scanned area. Using a computerized pattern generator and a new scanning algorithm, thermal heat is dispersed within the epidermis while evenly treating damaged areas. This particular algorithm dramatically reduces pain, the need for anesthesia, and recovery time. Single treatments can show a significant improvement in skin texture and color.

RISKS

Erythema (redness of skin): The laser-treated areas have a distinctive redness representing increased blood flow due to the growth of the superficial tissue and collagen. It will gradually fade during the first two days.

Inflammation (swelling): Treatment with the MiXto System presents a minimal amount of swelling that can last up to 24 hours.

Hyperpigmentation (tanning): This is presented after the third day of the procedure but then begins to shed and reveal younger healthy looking skin.

A personal interview and clinical examination will be conducted to obtain relevant facts about your medical and dermatologic history, and any medications you are currently taking or have taken in the recent past. If you are prone to cold sores, you need to ask your doctor for an antivirus prescription before the treatment.

Fractional Treatment with the MiXto SX has been proven safe for the face, neck, chest, and hands.

INSTRUCTIONS

Cool compresses on and off (i.e. 20 minutes to the eyes for the first 24 hours). Typically, take some gauze and put it into ice water and wring it out. You do not need to do this while asleep. In some patients who have had laser resurfacing, the cool compresses hurt the laser areas. The use of col compresses is optional.

Mild Pain Medicine: Ibuprofen or plain Tylenol for mild to moderate pain. Moderate to severe pain—Percocet ½ tablet every 3-4 hours with food. Laser areas on lower eyelids or full face should be treated topically with Aquaphor for three days, and switch to Hydraphase or Cetaphil lotion on day 4.

Antibiotic by mouth—none unless recommended by doctor

Continue Zovirax(Acyclovir) one tablet 3 times a day for 5 days after surgery (start two days before surgery) Total of 7 days. This is to prevent an outbreak of cold sores.

Zofran 8 mg (Ondansetron) sublingual every 8 hours ONLY IF NEEDED for nausea. Call Dr. Altman if you are having nausea.

Do not pick at any scabs that may form. Avoid sun or trauma to the area for about 1 week. Inform Dr. Altman if you develop any open sores or infections.

If you have severe pain, swelling, or loss of vision, please call the office at once. It is normal for some cloudiness of vision from the ointment. You can use refresh tears as needed to wash out your eyes, or to give moisture if dry.

Appointment
Call Dr. Altman if you have any problems or concerns.
Donald Altman, M.D.

Informed Consent for Fractional CO2 Laser Procedures

PLEASE INITIAL

I understand that Micro-Spot F vary depending on the condition o	ractional CO2 laser is intended skin re	ejuvenation and that results may
-	ne treatment may be needed for optil	mal results
	known conditions that could make tr	
 Pregnancy or currently bre 		
Herpes Simplex (cold sores	or fever blisters) in area to by treate	d
 History of keloid scarring 		
Use of medications that in-	crease photosensitivity	
 Tattoo in treated area 		
Auto-Immune Disorders		
 Diabetes 		
 Using Accutane 		
• Use of aspirin, Ibuprofen, a	and other anti-inflammatory medicati	ons
 UV light exposure 3 days p 	rior and 3 days post treatment	
I am fully aware of the possible	complications from the Micro-Spot F	Fractional CO2 laser, including
but not limited to:		
 Mild to moderate discomfe 	ort	
 Redness or swelling at the 	treated area	
 Sun sensitivity 		
 Skin sensitivity 		
 Blistering 		
 Bruising/Swelling/Infection 	n (rare)	
 Temporary or Permanent s 	skin pigmentation changes (rare): hyp	o- or hyperpigmentation
 Scarring (rare) 		
• Eye Exposure: Protective e	yewear will be provided to me	
I understand that there are oth	ner alternative treatments for skin reju	uvenation besides Micro-Spot
Fractional CO2 Laser.		
	be taken for my medical records	
	adult of at least 18 years of age and t	hat this consent is given freely
and voluntarily	1.1.1.2	
	closure and that it supersedes any production of the control of th	
	n to have any questions answered.	offilation and that i have had
samelent opportunity for alsoussie	The flave any questions answered.	
Signature of Patient/Guardian	Print Name/Relationship	 Date
Signature of Futienty Guardian	Thire Name, Relationship	Dutc
Signature of Witness	Print Name/Relationship	 Date